



Total and lean fish intake is positively associated with bone mineral density in older women in the community-based Hordaland Health Study

Hanne Rosendahl-Riise¹ · Therese Karlsson¹ · Christian A. Drevon² · Ellen M. Apalset^{3,4} · Ottar K. Nygård^{1,5} · Grethe S. Tell⁴ · Jutta Dierkes⁶

Received: 1 November 2017 / Accepted: 9 March 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Purpose Fish is a source of various nutrients beneficial for bone health, but few studies have investigated the association between bone mineral density (BMD) and fish consumption. Thus, the aim was to investigate the relationship between total fish intake and BMD and between both lean and fatty fish intake and BMD.

Method These cross-sectional analyses include 4656 participants in the Hordaland Health Study, a community-based study conducted in 1997–1999. The study includes two birth cohorts of men and women from Hordaland county (Norway) born in 1950–1951 and 1925–1927. BMD was measured by dual-energy X-ray absorptiometry and dietary intake was obtained from a semi-quantitative food-frequency questionnaire.

Results The average total fish intake was 33 ± 18 g/1000 kcal and was primarily lean fish. Older women had significantly lower BMD than older men and middle-aged men and women. In older women, total and lean fish intake (50 g/1000 kcal) was significantly and positively associated with BMD also after multivariate adjustments (β -coefficient 0.018, $p=0.017$ and 0.026, $p=0.021$).

Conclusion A high intake of fish, in particular lean fish, was positively associated with BMD in older women. No association between intake of fatty fish and BMD was found in either of the age and sex groups.

Keywords Diet · Food-frequency questionnaire · Fatty fish · Lean fish · Bone mineral density · Osteoporosis

Abbreviations

BMD Bone mineral density
DHA Docosahexaenoic acid

DPA Docosapentaenoic acid
DXA Dual-energy X-ray absorptiometry
EPA Eicosapentaenoic acid
FFQ Food-frequency questionnaire
HUSK Hordaland Health Study
NNR 2012 Nordic Nutritional Recommendation 2012
n3 PUFA Omega-3 polyunsaturated fatty acids

✉ Hanne Rosendahl-Riise
hanne.gjessing@uib.no

¹ Department of Clinical Science, Faculty of Medicine, University of Bergen, P.O. Box 7804, Laboratory Building 8th Floor, 5020 Bergen, Norway

² Department of Nutrition, Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, Oslo, Norway

³ Department of Rheumatology, Haukeland University Hospital, Bergen, Norway

⁴ Department of Global Public Health and Primary Care, University of Bergen, Bergen, Norway

⁵ Department of Heart Disease, Haukeland University Hospital, Bergen, Norway

⁶ Department of Clinical Medicine, Faculty of Medicine, University of Bergen, Bergen, Norway

Introduction

Osteoporosis is a major public health challenge, especially in an aging population, with the most severe consequence being fractures of the hip, wrist, or spine. The diagnosis of osteoporosis is made either after a low-energy fracture or by measurement of bone mineral density (BMD), preferably by dual-energy X-ray absorptiometry (DXA) technique. A BMD below 2.5 standard deviations of the average of young healthy adults is indicative for osteoporosis, applying age, and sex-specific cutoffs [1, 2]. In humans, BMD reaches

its peak in the third decade of life [3], and it decreases throughout life, with the fastest decline among women in the peri- and early postmenopausal state. This could be due to loss of endogenous estrogen, which in turn is associated with increased production of pro-inflammatory cytokines as mediators for the accelerated bone loss. It could also be due to other cell-autonomous age-related factors [4]. Men have higher BMD levels than women of similar age due to larger bones and thicker bone cortex [5].

Non-modifiable risk factors for low BMD include old age, female gender, and genetics, where genetics is proposed to predict 60–80% of the variability in bone mass [6]. Genetic risk factors include family history of osteoporosis [7, 8], ethnic differences in BMD [9, 10], and individual genetic variations [11]. Modifiable factors associated with low BMD are low lean body mass, alcohol consumption, smoking, physical inactivity, and use of osteoporosis inducing drugs like glucocorticoids. In recent years, diet has been given attention as a modifiable risk factor associated with bone health throughout life [6]. Although the role of specific nutrients or foods is debated, there is reasonable consensus that calcium, vitamin D [12, 13], vitamin K [14, 15], and perhaps n3 PUFA [16, 17] are important for bone health. Recently, the National Osteoporosis Foundation stated that there is moderate-to-strong evidence for calcium, vitamin D, and dairy consumption having a positive effect on peak bone mass [6]. The role of protein intake is a matter of debate, but recently, a high protein intake was reported to be associated with higher bone mass [18, 19]. Fish is a good source of nutrients associated with prevention of osteoporosis, such as high-quality protein, n3 PUFA, and vitamin D. Only a few studies have investigated the association between fish consumption and BMD. In the Framingham Osteoporosis Study, a protective effect of high intake of fish (≥ 3 servings/week) was found on bone loss [20]. In another large US prospective cohort of older adults, high fish consumption was associated with lower BMD [21]. Two Chinese studies found beneficial effects of fish intake on BMD and risk of osteoporosis [22, 23]. In addition, in Spanish premenopausal women, a positive association between fish intake and BMD was reported [24]. Due to these conflicting data, there is a need for more studies on the effect of fish intake in different populations with marked variations in both BMD and fish intake. Fish is a heterogeneous food group and populations with high total fish intake allow analysis of different types of fish. In general, Norwegians have a high intake of fish, with higher intake in older than in younger groups [25, 26]. Due to the habitually high fish intake in Norway, the present cohort is well suited to explore the association between BMD and overall fish intake, as well as lean and fatty fish intake. Thus, the main objective of this cross-sectional study was to investigate the relationship between intake of total, lean, and fatty fish and BMD in the Hordaland Health Study (HUSK).

Subjects and methods

Study population

The current work is a cross-sectional study of the large community-based Norwegian Hordaland Health Study (HUSK). HUSK was conducted from 1997 to 1999 as a collaboration between the University of Bergen, University of Oslo, local health services, and the Norwegian Institute of Public Health. Participants in HUSK are from two birth cohorts born either in 1925–1927 (older cohort) or in 1950–1951 (middle-aged cohort) from Hordaland county in Western Norway.

In 1997–1999, information on dietary intake and bone mineral density (BMD) was collected in about 4700 participants, allowing analysis of the association between dietary intake and BMD in both middle-aged and older men and women. More information about HUSK can be found at <http://husk-en.b.uib.no/>.

Dietary assessment

Habitual dietary intake (reflecting the previous year) was estimated using a 169-item FFQ developed at the Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo [27]. The questionnaire was handed out on the day of the health examination and then filled out at home. The questionnaire was later mailed to the HUSK Project Center in Bergen. Portion size was considered (e.g. slice, glass etc.) and questions on supplement use were included in the FFQ. Daily food (including fish consumption) and nutrient intakes were calculated using a food database and software system (Kostberegningssystem, version 3.2; University of Oslo, Norway). The FFQ has been compared against a weighted dietary record and fatty acid composition in serum phospholipids [28, 29]. For the dietary intake, an energy intake lower than 700 or 800 kcal and higher than 360 or 4200 kcal for women and men, respectively, was considered unreasonable and removed from the analysis, leaving 4656 participants with dietary records.

The questions related to dietary fish intake have been described in detail elsewhere [30]. Briefly, in addition to total fish (without shellfish), fish intake was divided into fatty fish (herring, mackerel, salmon, trout, and fish used as spread) and lean fish (cod, pollock, and haddock). The nutrient density method was used for energy adjustments [31] of all dietary variables and either stated as g/1000 kcal or percentage of total energy intake. In the multiple linear regression models, total fish, lean fish, and fatty fish were presented as 50 g/1000 kcal. The total

marine n3 PUFA intake was calculated by combining the variables for eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA), and docosahexaenoic acid (DHA).

Use of fish oil, cod liver oil (oil or capsules), calcium supplements, and vitamin D supplements was assessed in the FFQ. Participants who reported using such supplements more than once a week were defined as users. Alcohol intake was self-reported and was converted into g/day. Sex-specific cutoffs were used, and one unit of alcohol was defined as 10 g/day in accordance with the Nordic Nutrition Recommendations 2012 [32]. The intake was grouped into four categories; 0 = 0 g/day; 1 = women: > 0–10 g/day; men: > 0–20 g/day; 2 = women: > 10–20 g/day; men: > 20–30 g/day; 3 = women: > 20 g/day; men: > 30 g/day.

Clinical data

The HUSK measurements included a measurement of BMD of 5377 participants by DXA. The DXA measurement was performed at a different appointment after the initial visit to the Project Center. Measurements of BMD have been described in detail elsewhere [33]. Briefly, BMD of the femoral neck and total hip (g/cm^2) was measured by a DXA (EXPERT-XL; Lunar Company Inc, Madison, Wis, USA). The left hip was scanned unless there was a history of the previous fracture or surgery. The DXA measurements also allow calculation of body composition, that is fat mass and lean mass [34]. Weight and height were measured with the participants wearing light clothing without shoes, to the nearest 0.5 kg and 1 cm, respectively.

Covariate assessment

Self-administered questionnaires provided information regarding current estrogen therapy, physical activity (hard and light), and smoking (current/former/never smoked). Physical activity was categorized as by Vinknes et al. [35]. Categories for light physical activity were 0 (none), 0.25 (< 1 h/week), 0.5 (1–2 h/week), or 1.0 (≥ 3 h/week) and for hard physical activity were 0 (none), 0.5 (< 1 h/week), 1.0 (1–2 h/week), or 2.0 (≥ 3 h/week). The sum of these scores was calculated and used in the multivariate models. Smoking habits were categorized as current smoker, former smoker, and never smoker. In addition to the self-reported smoking habits, cotinine was measured as a marker of recent nicotine exposure. Cotinine was measured in EDTA plasma stored at -80 °C until analyzed at Bevital A/S (<http://www.bevital.no>), Bergen, Norway by LC/MS/MS. Smokers were defined due to cotinine levels ≥ 85 nmol/L [36, 37].

Statistical analysis

Continuous variables are presented as means and standard deviation and categorical variables as percentages. Differences between sex and age were assessed using Mann–Whitney *U* test for continuous variables and Fischer's exact test for categorical variables. Fish intake was also categorized into quartiles, calculated separately for each age group and sex. Differences in characteristics across quartiles of total fish intake were analyzed using linear regression for continuous variables and logistic regression for dichotomous variables.

The association of fish intake with BMD was analyzed both in quartiles of fish intake and fish intake as continuous variable. Multiple linear regression analyses were performed to assess the association between intake of either total, fatty, or lean fish and BMD (by age group and sex) with adjustment for potential confounders. Due to missing values in confounders (1.3–6.2%), the multivariate analysis included 4279 participants, whereas the energy-adjusted model included 4656 participants. The number of nonconsumers of lean and fatty fish was low, 405 (8.7%) and 255 (5.5%), respectively. The nonconsumers were categorized in quartile one.

The statistical software SPSS for Windows version 22 (IBM, NY, USA) was used for statistical analyses. A two-sided *p* value < 0.05 was considered statistically significant.

Results

The current analysis is based on 4656 participants from the HUSK study. Eligibility and selection of participants are shown in Fig. 1.

Characteristics

Characteristics of the study population, stratified by age group and sex, are presented in Table 1. There were more women (57%) than men in the total cohort. Femoral neck BMD and total hip BMD were higher in men than in women and higher in the middle-age cohorts than in the older.

A significantly higher proportion of both middle-aged and older women did not engage in hard physical activity in their leisure time, compared to men in the same age group. In the older individuals, men were more likely to be former smokers than women, while a higher proportion of women had never smoked. Significantly, more women than men reported no alcohol consumption. Men had lower fat mass and higher lean mass than women, and the older women had significantly more fat mass and lower lean mass than the




Fig. 1 Flowchart of the study population available for analysis from the Hordaland Health Study (HUSK). DXA dual-energy x-ray absorptiometry; FFQ food frequency questionnaire

middle-aged women. About one in six women used estrogen therapy.

Dietary intake by quartiles of fish intake

Fish intake was categorized into sex- and age-specific quartiles. Dietary intake for nutrients and food groups are presented across quartiles of fish intake in Table 2 (middle-aged and older men) and Table 3 (middle-aged and older women). Energy intake was not different between fish intake across the quartiles in any age or sex group. Older men had the highest total fish intake of all the age and sex groups. The intake of lean fish in older men was almost twice the intake in middle-aged men. There were also small, but significant differences in fatty fish intake. Age group differences in the female cohorts were less pronounced, but the older women had higher total fish intake and higher lean fish intake than the middle-aged women.

Dairy intake was similar in all the quartiles of fish intake in men, but among women, high fish consumers had low consumption of milk products. For all groups, high fish

consumers usually had a higher meat intake than low fish consumers. High fish consumers also had a considerably higher vegetable intake than the low fish consumers. There was no difference in fruit and berries consumption across the quartiles among men or women nor between the two age groups.

In the total cohort, 36% used cod liver oil weekly, with the older men having the highest intake (40%). Supplementation of fish oil and vitamin D was not common. In the older women, 13% used calcium supplements, which is considerably higher than in the other groups.

Intake of vitamin D and n3 PUFA from food and supplements, and protein intake (both as energy percent and as g per kg body weight) increased across quartiles of fish intake in all cohorts. There was no difference in total calcium intake across quartiles.

Fish intake and bone mineral density

The Spearman's rho between total hip BMD and femoral neck BMD was in middle-aged men: $r=0.90$, older men: $r=0.89$, middle-aged women: $r=0.88$ and older women: $r=0.86$. Use of total hip BMD as outcome variable did not change the results substantially (data not shown). Femoral neck BMD increased across quartiles of fish intake in middle-aged men and older women, while there was no significant association in middle-aged women or older men (Fig. 2).

The linear regression analysis on the association of fish intake with femoral neck BMD is presented in Table 4. In middle-aged men and older women, total fish intake was positively associated with BMD (Model 1; adjusted for total energy intake). After additional adjustments for BMI, physical activity score, cotinine > 85 nmol/L, and alcohol consumption (Model 2), the association was no longer significant for middle-aged men, but remained significant in the older women. Further analysis of the type of fish revealed that high lean fish intake was significantly associated with high femoral neck BMD in older women in the fully adjusted model, whereas fatty fish did not show such an association. A sensitivity analysis leaving out the nonconsumers, did not change the results (data not shown).

Discussion

In this population-based study in Norwegians with high habitual fish intake, a positive significant association between total fish intake and BMD was observed in older women (70–74 years), but not in older men or in the middle-aged cohorts (46–49 years). The effect remained stable even after adjustment for various covariates known to be associated with BMD. The association of total fish intake with

Table 1 Characteristics of 4656 middle-aged (46–49 years) and older (70–74 years) men and women in the Hordaland Health Study

	Total cohort	Middle-aged		Older	
		Men (<i>n</i> = 1052)	Women (<i>n</i> = 1605)	Men (<i>n</i> = 962)	Women (<i>n</i> = 1037)
Age (years)		47 ± 1	47 ± 1	72 ± 1	72 ± 1
Women (%)	56.7				
Weight (kg) ^{a,c,d,e}	74.1 ± 13.4	84.0 ± 11.9	68.3 ± 1.6	79.7 ± 11.1	67.6 ± 11.2
Height (m) ^{a,c,d,e,f}	1.70 ± 0.09	1.79 ± 0.06	1.66 ± 0.06	1.75 ± 0.06	1.61 ± 0.05
BMI (kg/m ²) ^{a,c,f}	25.7 ± 3.8	26.2 ± 3.3	24.8 ± 4.0	26.0 ± 3.2	26.2 ± 4.2
Fat mass (kg) ^{a,c,d,e,f}	23.4 ± 9.6	20.5 ± 9.1	24.4 ± 9.6	21.3 ± 8.6	26.9 ± 9.5
Lean mass (kg) ^{a,c,d,e,f}	47.2 ± 10.6	59.9 ± 6.2	40.3 ± 4.5	55.1 ± 5.8	37.7 ± 4.3
Femoral neck BMD (g/cm ²) ^{c,d,e,f}	0.911 ± 0.15	1.000 ± 0.14	0.961 ± 0.12	0.901 ± 0.12	0.763 ± 0.11
Total hip BMD (g/cm ²) ^{c,d,e,f}	0.950 ± 0.16	1.031 ± 0.14	0.986 ± 0.13	0.962 ± 0.15	0.800 ± 0.12
Hard physical activity (%) ^a					
None ^{c,d,e,f}	3.5	2.4	4.2	2.4	5.6
<1 h/week ^{c,d,e,f}	22.2	20.4	18.5	19.5	32.7
1–2 h/week ^{d,e}	45.9	45.2	45.1	47.6	44.5
≥3 h/week ^{c,d,f}	28.4	32.0	32.2	30.4	17.2
Smoking habits (%)					
Current smoker ^{d,e,f}	27.4	35.4	36.3	17.8	14.5
Former smoker ^{c,d,e}	35.2	33.4	26.6	60.7	26.8
Never smoked ^{c,d,e,f}	40.0	35.6	40.0	24.6	60.4
Cotinine ≥ 85 nmol/L (%) ^{a,c,f}	27.4	35.3	36.4	17.7	14.5
Alcohol categories (%) ^b					
None ^{c,d,e,f}	27.5	9.1	20.6	29.8	53.5
Low ^{c,d,e,f}	63.5	80.0	67.7	62.2	41.5
Moderate ^{c,e,f}	6.7	6.7	10.0	4.0	4.3
High ^{s,d,f}	2.7	4.2	2.3	4.1	0.7
Current estrogen therapy (for women) (%) ^f	16.8	NA	17.9	NA	15.1

Values represented mean ± SD or %. *p* Values for the difference between the age and sex groups were calculated using Mann–Whitney *U* test or Fischer exact test

^a*n* = 4279–4653

^bNone: 0 g/day, low: women: >0–10 g, men: >0–20 g, moderate: women: >10–20 g, men: >20–30 g, high: women: >20 g, men: >30 g

^cSignificant (*p* ≤ 0.05) difference between the middle-age by sex

^dSignificant (*p* ≤ 0.05) difference between the older by sex

^eSignificant (*p* ≤ 0.05) difference between the men by age

^fSignificant (*p* ≤ 0.05) difference between the women by age

BMD was mainly due to the high intake of lean fish, whereas fatty fish intake was not significantly associated with BMD in any of the cohorts.

Comparison with other studies

Although several studies have focused on fish intake and BMD, these studies are difficult to compare due to differences in age range, sex, and ethnicity. There are also a variety in amount and type of fish consumed, and how fish intake is monitored. In the present study, the association of fish intake with BMD was evaluated from dietary intake data obtained from an FFQ, energy-adjusted and analyzed as a single food group, but with separate analyses for fatty and

lean fish. A similar approach has been used in predominantly Caucasian populations from the US and Europe [20, 21, 24, 38] and in Asian cohorts [22, 23, 38, 39]. With the exception of the study by Virtanen et al. [21], these studies report a positive association of fish intake and BMD, despite differences in fish classification and assessment of dietary fish intake. However, the association was only significant in the postmenopausal Chinese women with very high fish intake [22], in old rural Chinese women with fish intake > 250 g/week [23], in the high consuming Spanish premenopausal women [24], and in Koreans reporting high fish intake [38]. The latter study showed that many other factors were important for BMD in old adults, as despite high fish intake and positive associations with BMD in the Koreans, the absolute

Table 2 Dietary characteristics and measures of BMD in 1052 middle-aged (46–49 years) and 962 older (70–74 years) men by quartiles of total fish intake in the Hordaland Health Study

Total fish, g (min–max)	Middle-aged men				Older men				<i>p</i> for trend		
	Total	1st quartile (<i>n</i> = 263) 1.1–20.9	2nd quartile (<i>n</i> = 263) 20.9–30.6	3rd quartile (<i>n</i> = 263) 30.6–42.5	4th quartile (<i>n</i> = 263) 42.5–162.7	Total	1st quartile (<i>n</i> = 240) 3.0–33.0	2nd quartile (<i>n</i> = 241) 33.1–46.7		3rd quartile (<i>n</i> = 241) 46.7–64.1	4th quartile (<i>n</i> = 240) 64.3–186.3
Total energy (kcal)	2533 ± 697	2487 ± 752	2621 ± 654	2527 ± 608	2496 ± 759	2082 ± 573	2062 ± 559	2125 ± 602	2140 ± 579	1999 ± 543	0.291
Protein (E%)	15.7 ± 2.2	14.5 ± 2.1	15.2 ± 1.7	16.0 ± 1.8	17.3 ± 2.0	16.1 ± 2.3	14.4 ± 1.9	15.6 ± 1.8	16.3 ± 1.7	18.0 ± 2.1	<0.001
n-3 long chained PUFA (E%) ^a	0.07 ± 0.07	0.04 ± 0.06	0.05 ± 0.04	0.07 ± 0.06	0.1 ± 0.09	0.11 ± 0.10	0.06 ± 0.06	0.09 ± 0.09	0.11 ± 0.08	0.16 ± 0.11	<0.001
Protein g/kg body weight	1.2 ± 0.4	1.1 ± 0.4	1.2 ± 0.3	1.2 ± 0.3	1.3 ± 0.4	1.1 ± 0.3	1.0 ± 0.3	1.0 ± 0.3	1.1 ± 0.3	1.1 ± 0.4	<0.001
Vitamin D (g/1000 kcal) ^b	4 ± 3	4 ± 3	4 ± 3	4 ± 3	6 ± 4	6 ± 4	4 ± 3	5 ± 4	6 ± 4	7 ± 5	<0.001
Calcium (g/1000 kcal) ^b	384 ± 120	392 ± 133	378 ± 119	386 ± 111	378 ± 115	377 ± 119	451 ± 136	390 ± 137	385 ± 123	369 ± 101	0.004
Supplements (%)											
Fish oil	8.4	7.2	4.9	9.1	12.2	5.3	3.8	5.4	4.6	7.5	0.304
Cod liver oil	38.9	35.7	38.9	38.4	42.6	40.1	30.8	43.2	46.5	40.0	0.004
Vitamin D	1.5	0	2.7	1.5	1.9	1.9	0.8	3.3	0.8	2.5	0.148
Calcium	1.4	0.8	1.9	1.1	1.9	2.8	2.1	2.5	2.9	3.8	0.725
Food intake (g/1000 kcal)											
Total fish	33 ± 18	13 ± 5	26 ± 3	36 ± 3	58 ± 16	51 ± 25	23 ± 7	40 ± 4	55 ± 5	85 ± 20	<0.001
Lean fish	12 ± 10	4 ± 4	9 ± 5	14 ± 7	21 ± 13	23 ± 17	9 ± 7	17 ± 8	24 ± 10	40 ± 19	<0.001
Fatty fish	10 ± 10	2 ± 2	4 ± 4	6 ± 5	11 ± 9	15 ± 13	3 ± 4	6 ± 6	9 ± 7	16 ± 13	<0.001
Vegetables	79 ± 61	66 ± 58	73 ± 59	83 ± 57	95 ± 66	91 ± 59	79 ± 54	88 ± 60	91 ± 50	106 ± 69	<0.001
Fruit and berries	97 ± 65	94 ± 68	95 ± 59	95 ± 63	105 ± 68	116 ± 72	112 ± 73	117 ± 71	115 ± 67	118 ± 78	0.408
Meat	58 ± 23	56 ± 26	57 ± 21	59 ± 21	60 ± 23	45 ± 21	41 ± 21	46 ± 21	49 ± 20	46 ± 22	0.004
Milk products	156 ± 106	158 ± 118	157 ± 104	156 ± 101	155 ± 103	162 ± 105	167 ± 114	172 ± 107	159 ± 95	149 ± 104	0.031
Egg	7 ± 5	7 ± 6	7 ± 4	7 ± 5	8 ± 5	9 ± 7	8 ± 7	10 ± 8	8 ± 5	9 ± 6	0.793

The values represented mean ± SD or %. *P* for trend: linear regression for continuous variables and logistic regression for dichotomous variables

^aSum of eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA), and docosahexaenoic acid (DHA)

^bTotal intake from diet and supplements

Table 3 Dietary characteristics of 1605 middle-aged (46–49 years) and 1037 older (70–73 years) women by quartiles of total fish intake in the Hordaland Health Study

Total fish, g (min–max)	Middle-aged women				Older women				<i>p</i> for trend		
	Total	1st quartile (n=401)	2nd quartile (n=401)	3rd quartile (n=402)	4th quartile (n=401)	Total	1st quartile (n=259)	2nd quartile (n=259)		3rd quartile (n=260)	4th quartile (n=259)
Total energy (kcal)	1914±566	1894±552	1955±542	1891±482	1918±672	1627±497	1572±505	1628±483	1673±487	1636±510	0.085
Protein (E%)	16.4±2.4	15.1±2.2	15.7±1.9	16.6±1.9	18.1±2.3	<0.001	16.5±2.6	15.9±2.0	16.8±2.0	18.4±2.5	<0.001
Protein g/kg body weight	1.2±0.4	1.1±0.4	1.1±0.4	1.1±0.3	1.3±0.5	<0.001	1.0±0.4	0.9±0.4	1.0±0.3	1.1±0.4	<0.001
n-3 long chained PUFA (E%) ^a	0.09±0.10	0.06±0.08	0.07±0.08	0.09±0.10	0.12±0.13	<0.001	0.09±0.12	0.10±0.10	0.11±0.10	0.17±0.13	<0.001
Vitamin D (g/1000 kcal) ^b	5±4	4±4	4±4	5±4	6±4	<0.001	5±4	5±4	5±4	7±4	<0.001
Calcium (g/1000 kcal) ^b	409±127	421±136	402±116	410±116	405±137	0.395	451±136	470±153	443±126	432±130	0.342
Supplements (%)											
Fish oil	8.2	6.8	7.2	8.2	10.5	0.227	7.2	4.6	6.2	12.7	<0.001
Cod liver oil	34.0	30.0	36.2	36.3	33.7	0.145	32.9	34.0	29.3	35.5	0.489
Vitamin D	3.4	2.3	2.5	4.2	4.7	0.192	3.0	3.5	2.7	3.5	0.825
Calcium	7.5	5.3	8.0	9.2	7.7	0.199	12.7	10.0	13.5	14.3	0.493
Food intake (g/1000 kcal)											
Total fish	35±19	15±5	27±3	38±4	61±17	<0.001	51±25	18±6	35±4	78±21	<0.001
Lean fish	14±11	5±4	10±5	15±7	25±14	<0.001	23±17	7±5	15±7	38±20	<0.001
Fatty fish	10±10	3±3	4±4	6±5	12±10	<0.001	15±13	3±4	5±5	14±12	<0.001
Vegetables	121±77	102±74	112±73	131±78	141±78	<0.001	118±75	96±74	119±71	122±66	<0.001
Fruit and berries	135±83	133±94	136±81	137±78	135±80	0.736	152±102	142±108	162±109	155±86	0.532
Meat	56±23	54±24	56±22	57±22	58±24	0.018	40±20	34±18	40±19	42±20	<0.001
Milk products	134±99	140±112	136±94	137±96	124±93	0.033	192±117	216±136	204±116	162±99	<0.001
Egg	9±6	8±6	8±5	8±6	9±6	0.006	10±7	9±7	10±8	10±8	0.377

The values represented mean ± SD or %, *p* for trend: linear regression for continuous variables and logistic regression for dichotomous variables

^aSum of eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA), and docosahexaenoic acid (DHA)

^bTotal intake from diet and supplements




Fig. 2 Femoral neck BMD (g/cm^2) by quartiles (Q1–Q4) of total fish intake in middle-aged (46–49 years) and older (70–74 years) men and women in the Hordaland Health Study

measured BMD was higher in the Americans. This demonstrates the complexity of investigating the association between foods and BMD.

Whereas most studies have a cross-sectional design, there are two studies investigating fish intake and bone loss over time [20, 39]. These studies reported a protective effect of high fish consumption on bone loss, with significant

associations for dark fish and tuna in the Framingham study [20], and total fish intake in the Chinese study [39]. Longitudinal studies on the association of fish consumption and BMD may be even more relevant, as bone mass changes throughout life. Thus, nutritional intake including low intake of fish can be regarded as a modifiable risk factors for osteoporosis.

The association between fish intake and BMD has also been analyzed applying a dietary pattern approach. In these analyses, fish consumption is regarded as part of a healthy dietary pattern and was associated with BMD in Australian men and women > 50 years [40] and with Japanese premenopausal women [41], but not in older women in a Finnish study [42].

Possible mechanisms

Fish is a source of nutrients that have been associated with higher BMD, in particular protein [19], n3 PUFAs [16] and vitamin D [43]. A recent meta-analysis and systematic review showed that a higher protein intake was associated with higher BMD at most bone sites (albeit non-significant at most sites) and less bone loss over time [19]. However, it is difficult to translate this into dietary recommendations as ‘higher protein’ intake was defined differently in the studies included. The assumption that protein intake is important

Table 4 Unstandardized B coefficient for femoral neck BMD by intake of total fish, lean fish, and fatty fish obtained by multiple linear regression models in the middle-aged (46–49 years) and older (70–74 years) men and women in the Hordaland Health Study

	Model 1 ^a		Model 2 ^b	
	Unstandardized B coefficient (95% CI)	<i>p</i>	Unstandardized B coefficient (95% CI)	<i>p</i>
Middle-aged				
Men (<i>n</i> = 1052)				
Total fish (50 g/1000 kcal)	0.029 (0.007, 0.051)	0.011	0.006 (−0.016, 0.027)	0.612
Lean fish (50 g/1000 kcal)	0.035 (−0.007, 0.076)	0.099	0.007 (−0.033, 0.048)	0.714
Fatty fish (50 g/1000 kcal)	0.043 (0.003, 0.082)	0.033	0.013 (−0.025, 0.051)	0.386
Women (<i>n</i> = 1605)				
Total fish (50 g/1000 kcal)	0.010 (−0.006, 0.026)	0.204	0.006 (−0.010, 0.021)	0.475
Lean fish (50 g/1000 kcal)	−0.006 (−0.033, 0.022)	0.680	−0.009 (−0.036, 0.017)	0.484
Fatty fish (50 g/1000 kcal)	0.022 (−0.008, 0.052)	0.144	0.017 (−0.011, 0.046)	0.236
Older				
Men (<i>n</i> = 962)				
Total fish (50 g/1000 kcal)	0.015 (−0.003, 0.003)	0.094	0.002 (−0.016, 0.020)	0.842
Lean fish (50 g/1000 kcal)	0.010 (−0.017, 0.037)	0.460	−0.005 (−0.032, 0.022)	0.703
Fatty fish (50 g/1000 kcal)	0.007 (−0.028, 0.043)	0.680	−0.001 (−0.037, 0.035)	0.529
Women (<i>n</i> = 1037)				
Total fish (50 g/1000 kcal)	0.018 (0.004, 0.032)	0.014	0.018 (0.003, 0.032)	0.017
Lean fish (50 g/1000 kcal)	0.018 (−0.003, 0.039)	0.100	0.026 (0.004, 0.048)	0.021
Fatty fish (50 g/1000 kcal)	0.024 (−0.002, 0.051)	0.072	0.014 (−0.014, 0.042)	0.337

^aAdjusted for total energy intake

^bAdjusted for total energy intake (cont.), BMI (cont.), physical activity score (none/<1 h/week/1–2 h/week/≥ 3 h/week), cotinine > 85 nmol/L (yes/no), alcohol consumption (none/low/moderate/high)

for maintaining or even increasing BMD is based on the fact that 50% of bone tissue is made up of proteins [44]. Despite an average protein intake in the present study in accordance with the recommendations, about one-third of the older women consumed less than 0.8 g protein/kg body weight per day (Table 2), which is the recommended amount of protein in the Nordic nutritional recommendations 2012 (NNR 2012) [32]. However, even NNR 2012 states that this needs more investigation, and other societies such as the European Society for Clinical Nutrition and Metabolism [45] have recommended higher protein intake for older persons. Fish is a good source of protein and may contribute to a high protein intake. Historically, a high protein intake was thought to be associated with increased urinary loss of calcium [46]. However, current knowledge does not support an adverse effect of diets with high intake of protein as long as the intake of calcium is adequate [19]. It is also important to have a sufficient energy intake to prevent catabolism of body protein for energy purposes.

The positive association of fish consumption with BMD may be due to the n3 PUFA content of fish [17, 20, 21]. Proposed mechanisms explaining the association between n3 PUFA and BMD include the promotion of calcium absorption from the intestine, effect on osteoblastogenesis and osteoblast activity, reduction in inflammation, and modulation of peroxisome proliferators-activated receptor γ [16, 17]. Participants in HUSK consumed about twice the amount of n3 PUFA compared to participants in the previous studies [20, 21] due to fish intake, cod liver oil, and fish oil.

Surprisingly, only lean fish was associated with BMD among the older women. No association was found with fatty fish. Fatty fish and cod liver oil are important dietary sources of vitamin D, which is essential for the intestinal absorption of calcium, and a key factor in maintaining the calcium/phosphate homeostasis in serum. An association between high vitamin D status and high BMD has been reported [47]. Although fish is an important dietary source of vitamin D, synthesis of vitamin D in the skin is probably more essential [48]. However, skin synthesis of vitamin D in Norway at 60° northern latitude is only sufficient during the light period of the year. Thus, many Norwegians rely on nutrient sources of vitamin D a large part of the year. A recent meta-analysis concluded that consumption of a recommended amount of fish (300–400 g per week) was not enough to obtain a sufficient vitamin D status (50 nmol/L of 25OHD) [49]. However, due to the habitual intake of both cod liver oil and fatty fish, Norwegians on average have higher vitamin D intake than most other European populations [50].

Strength and limitations

The strength of this community-based study is the large sample size and inclusion of both women and men at different

ages. The study included only Caucasians, a population with a high risk of osteoporosis and fractures. Dietary intake was measured using a validated semi-quantitative FFQ [28]. The fish intake reported in our study was similar to intake data obtained in Norwegian dietary surveys [26], suggesting that the estimate was valid. Due to the habitually high fish consumption, it was possible to investigate lean and fatty fish separately, which has not been done by others. The assessment of several other factors associated with BMD allowed extensive adjustment in the multiple linear regression models.

The cross-sectional design of the analysis is a limitation due to non-causal explanation. In addition, we only had information on dietary intake the year before recruitment, whereas bone mass changes take place throughout a longer period of time. However, dietary habits seldom change rapidly, and the intakes reported are likely to represent habits extended throughout a longer period of life.

Conclusion

The total fish intake, in particular lean fish, was positively associated with BMD in older women. No association was found between fatty fish and BMD. Thus, amount and type of fish consumed should be investigated further in different populations as a possible modifiable factor for BMD.

Acknowledgements The Norwegian Seafood Research Fund (FHF) supported this work through fellowship funding the primary author. The funder had no role in the study design, analysis or interpretation of data, or in the writing of the manuscript.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval HUSK was performed according to the declaration of Helsinki and all participants provided written informed consent. The study protocol was approved by the Data Inspectorate and the Regional Committee for Medical Research Ethics.

References

1. WHO Study Group (1994) Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. WHO Technical Report Series
2. Kanis JA, Melton LJ 3rd, Christiansen C, Johnston CC, Khaltaev N (1994) The diagnosis of osteoporosis. *J Bone Miner Res* 9(8):1137–1141. <https://doi.org/10.1002/jbmr.5650090802>
3. Berger C, Goltzman D, Langsetmo L, Joseph L, Jackson S, Kreiger N, Tenenhouse A, Davison KS, Josse RG, Prior JC, Hanley DA (2010) Peak bone mass from longitudinal data: implications for the prevalence, pathophysiology, and diagnosis of osteoporosis.

- J Bone Miner Res 25(9):1948–1957. <https://doi.org/10.1002/jbmr.95>
4. Khosla S, Melton LJ, Riggs BL (2011) The unitary model for estrogen deficiency and the pathogenesis of osteoporosis: is a revision needed? *J Bone Miner Res* 26(3):441–451. <https://doi.org/10.1002/jbmr.262>
 5. Alswat KA (2017) Gender disparities in osteoporosis. *J Clin Med Res* 9(5):382–387. <https://doi.org/10.14740/jocmr2970w>
 6. Weaver CM, Gordon CM, Janz KF, Kalkwarf HJ, Lappe JM, Lewis R, O’Karma M, Wallace TC, Zemel BS (2016) The National Osteoporosis Foundation’s position statement on peak bone mass development and lifestyle factors: a systematic review and implementation recommendations. *Osteoporos Int*. <https://doi.org/10.1007/s00198-015-3440-3>
 7. Seeman E, Hopper JL, Bach LA, Cooper ME, Parkinson E, McKay J, Jerums G (1989) Reduced bone mass in daughters of women with osteoporosis. *N Engl J Med* 320(9):554–558. <https://doi.org/10.1056/nejm198903023200903>
 8. Soroko SB, Barrett-Connor E, Edelstein SL, Kritiz-Silverstein D (1994) Family history of osteoporosis and bone mineral density at the axial skeleton: the Rancho Bernardo Study. *J Bone Miner Res* 9(6):761–769. <https://doi.org/10.1002/jbmr.5650090602>
 9. Weaver CM, McCabe LD, McCabe GP, Novotny R, Van Loan M, Going S, Matkovic V, Boushey C, Savaiano DA (2007) Bone mineral and predictors of bone mass in white, Hispanic, and Asian early pubertal girls. *Calcif Tissue Int* 81(5):352–363. <https://doi.org/10.1007/s00223-007-9074-5>
 10. Bhudhikanok GS, Wang MC, Eckert K, Matkin C, Marcus R, Bachrach LK (1996) Differences in bone mineral in young Asian and Caucasian Americans may reflect differences in bone size. *J Bone Miner Res* 11(10):1545–1556. <https://doi.org/10.1002/jbmr.5650111023>
 11. Estrada K, Styrkarsdottir U, Evangelou E, Hsu YH, Duncan EL, Ntzani EE, Oei L, Albagha OM, Amin N, Kemp JP, Koller DL, Li G, Liu CT, Minster RL, Moayyeri A, Vandenput L, Willner D, Xiao SM, Yerges-Armstrong LM, Zheng HF, Alonso N, Eriksson J, Kammerer CM, Kaptoge SK, Leo PJ, Thorleifsson G, Wilson SG, Wilson JF, Aalto V, Alen M, Aragaki AK, Aspelund T, Center JR, Dailiana Z, Duggan DJ, Garcia M, Garcia-Giralt N, Giroux S, Hallmans G, Hocking LJ, Husted LB, Jameson KA, Khusainova R, Kim GS, Kooperberg C, Koromila T, Kruk M, Laaksonen M, Lacroix AZ, Lee SH, Leung PC, Lewis JR, Masi L, Mencej-Bedrac S, Nguyen TV, Nogue X, Patel MS, Prezelj J, Rose LM, Scollen S, Siggeirsdottir K, Smith AV, Svensson O, Trompet S, Trummer O, van Schoor NM, Woo J, Zhu K, Balcels S, Brandi ML, Buckley BM, Cheng S, Christiansen C, Cooper C, Dedoussis G, Ford I, Frost M, Goltzman D, Gonzalez-Macias J, Kahonen M, Karlsson M, Khusnutdinova E, Koh JM, Kollia P, Langdahl BL, Leslie WD, Lips P, Ljunggren O, Lorenc RS, Marc J, Mellstrom D, Obermayer-Pietsch B, Olmos JM, Petteersson-Kymmer U, Reid DM, Riancho JA, Ridker PM, Rousseau F, Slagboom PE, Tang NL, Urreizti R, Van Hul W, Viikari J, Zarrabeitia MT, Aulchenko YS, Castano-Betancourt M, Grundberg E, Herrera L, Ingvarsson T, Johannsdottir H, Kwan T, Li R, Luben R, Medina-Gomez C, Palsson ST, Reppe S, Rotter JJ, Sigurdsson G, van Meurs JB, Verlaan D, Williams FM, Wood AR, Zhou Y, Gautvik KM, Pastinen T, Raychaudhuri S, Cauley JA, Chasman DI, Clark GR, Cummings SR, Danoy P, Dennison EM, Eastell R, Eisman JA, Gudnason V, Hofman A, Jackson RD, Jones G, Jukema JW, Khaw KT, Lehtimaki T, Liu Y, Lorentzon M, McCloskey E, Mitchell BD, Nandakumar K, Nicholson GC, Oostra BA, Peacock M, Pols HA, Prince RL, Raitakari O, Reid IR, Robbins J, Sambrook PN, Sham PC, Shuldiner AR, Tylavsky FA, van Duijn CM, Wareham NJ, Cupples LA, Econs MJ, Evans DM, Harris TB, Kung AW, Psaty BM, Reeve J, Spector TD, Streeten EA, Zillikens MC, Thorsteinsdottir U, Ohlsson C, Karasik D, Richards JB, Brown MA, Stefansson K, Uitterlinden AG, Ralston SH, Ioannidis JP, Kiel DP, Rivadeneira F (2012) Genome-wide meta-analysis identifies 56 bone mineral density loci and reveals 14 loci associated with risk of fracture. *Nat Genet* 44(5):491–501. <https://doi.org/10.1038/ng.2249>
 12. Tang BM, Eslick GD, Nowson C, Smith C, Bensoussan A (2007) Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet* 370(9588):657–666. [https://doi.org/10.1016/s0140-6736\(07\)61342-7](https://doi.org/10.1016/s0140-6736(07)61342-7)
 13. Holvik K, Ahmed LA, Forsmo S, Gjesdal CG, Grimnes G, Samuelsen SO, Schei B, Blomhoff R, Tell GS, Meyer HE (2013) Low serum levels of 25-hydroxyvitamin D predict hip fracture in the elderly: a NOREPOS study. *J Clin Endocrinol Metab* 98(8):3341–3350. <https://doi.org/10.1210/jc.2013-1468>
 14. Booth SL, Tucker KL, Chen H, Hannan MT, Gagnon DR, Cupples LA, Wilson PW, Ordovas J, Schaefer EJ, Dawson-Hughes B, Kiel DP (2000) Dietary vitamin K intakes are associated with hip fracture but not with bone mineral density in elderly men and women. *Am J Clin Nutr* 71(5):1201–1208
 15. Finnes TE, Lofthus CM, Meyer HE, Sogaard AJ, Tell GS, Apalset EM, Gjesdal C, Grimnes G, Schei B, Blomhoff R, Samuelsen SO, Holvik K (2015) A combination of low serum concentrations of vitamins K and D is associated with increased risk of hip fractures in elderly Norwegians: a NOREPOS study. *Osteoporos Int*. <https://doi.org/10.1007/s00198-015-3435-0>
 16. Maggio M, Artoni A, Lauretani F, Borghi L, Nouvenne A, Valenti G, Ceda GP (2009) The impact of omega-3 fatty acids on osteoporosis. *Curr Pharm Des* 15(36):4157–4164
 17. Orchard TS, Pan X, Cheek F, Ing SW, Jackson RD (2012) A systematic review of omega-3 fatty acids and osteoporosis. *Br J Nutr* 107(Suppl 2):S253–S260. <https://doi.org/10.1017/s0007114512001638>
 18. Darling AL, Millward DJ, Torgerson DJ, Hewitt CE, Lanham-New SA (2009) Dietary protein and bone health: a systematic review and meta-analysis. *Am J Clin Nutr* 90(6):1674–1692. <https://doi.org/10.3945/ajcn.2009.27799>
 19. Shams-White MM, Chung M, Du M, Fu Z, Insogna KL, Karlens MC, LeBoff MS, Shapses SA, Sackey J, Wallace TC, Weaver CM (2017) Dietary protein and bone health: a systematic review and meta-analysis from the National Osteoporosis Foundation. *Am J Clin Nutr*. <https://doi.org/10.3945/ajcn.116.145110>
 20. Farina EK, Kiel DP, Roubenoff R, Schaefer EJ, Cupples LA, Tucker KL (2011) Protective effects of fish intake and interactive effects of long-chain polyunsaturated fatty acid intakes on hip bone mineral density in older adults: the Framingham Osteoporosis Study. *Am J Clin Nutr* 93(5):1142–1151. <https://doi.org/10.3945/ajcn.110.005926>
 21. Virtanen JK, Mozaffarian D, Cauley JA, Mukamal KJ, Robbins J, Siscovick DS (2010) Fish consumption, bone mineral density, and risk of hip fracture among older adults: the cardiovascular health study. *J Bone Miner Res* 25(9):1972–1979. <https://doi.org/10.1002/jbmr.87>
 22. Chen YM, Ho SC, Lam SS (2010) Higher sea fish intake is associated with greater bone mass and lower osteoporosis risk in postmenopausal Chinese women. *Osteoporos Int* 21(6):939–946. <https://doi.org/10.1007/s00198-009-1029-4>
 23. Zalloua PA, Hsu YH, Terwedow H, Zang T, Wu D, Tang G, Li Z, Hong X, Azar ST, Wang B, Boussein ML, Brain J, Cummings SR, Rosen CJ, Xu X (2007) Impact of seafood and fruit consumption on bone mineral density. *Maturitas* 56(1):1–11. <https://doi.org/10.1016/j.maturitas.2006.05.001>
 24. Calderon-Garcia JF, Moran JM, Roncero-Martin R, Rey-Sanchez P, Rodriguez-Velasco FJ, Pedrera-Zamorano JD (2012) Dietary habits, nutrients and bone mass in Spanish premenopausal

- women: the contribution of fish to better bone health. *Nutrients* 5(1):10–22. <https://doi.org/10.3390/nu5010010>
25. Totland TH, Melnæs BK, Lundberg-Hallén N, Helland-Kigen KM, Lund-Blix NA, Myhre JB, Johansen AMW, Løken EB, Andersen LF (2012) Norkost 3—a National Dietary Survey in among adults aged 18–70 years in Norway, 2010–2011
 26. Johansson L, Solvoll K (1999) Norkost 1997—Landsomfattende kostholdsundersøkelse blant menn og kvinner i alderen 16–79 år. Statens råd for ernæring og fysisk aktivitet
 27. Nes M, Frost Andersen L, Solvoll K, Sandstad B, Hustvedt BE, Lovo A, Dreven CA (1992) Accuracy of a quantitative food frequency questionnaire applied in elderly Norwegian women. *Eur J Clin Nutr* 46(11):809–821
 28. Andersen LF, Solvoll K, Johansson LR, Salminen I, Aro A, Dreven CA (1999) Evaluation of a food frequency questionnaire with weighed records, fatty acids, and alpha-tocopherol in adipose tissue and serum. *Am J Epidemiol* 150(1):75–87
 29. Andersen LF, Solvoll K, Dreven CA (1996) Very-long-chain n-3 fatty acids as biomarkers for intake of fish and n-3 fatty acid concentrates. *Am J Clin Nutr* 64(3):305–311
 30. Nurk E, Dreven CA, Refsum H, Solvoll K, Vollset SE, Nygard O, Nygaard HA, Engedal K, Tell GS, Smith AD (2007) Cognitive performance among the elderly and dietary fish intake: the Hordaland Health Study. *Am J Clin Nutr* 86(5):1470–1478
 31. Willett WC, Howe GR, Kushi LH (1997) Adjustment for total energy intake in epidemiologic studies. *Am J Clin Nutr* 65 (4 Suppl):1220S–1228S (**discussion 1229S–1231S**)
 32. Nordic Council of Ministers 2014 (2012) Nordic Nutritional Recommendations 2012—integrating nutrition and physical activity, 5th edn. Nordic Council of Ministers, Copenhagen
 33. Gjesdal CG, Aanderud SJ, Haga HJ, Brun JG, Tell GS (2004) Femoral and whole-body bone mineral density in middle-aged and older Norwegian men and women: suitability of the reference values. *Osteoporos Int* 15(7):525–534. <https://doi.org/10.1007/s00198-003-1573-2>
 34. Gjesdal CG, Halse JI, Eide GE, Brun JG, Tell GS (2008) Impact of lean mass and fat mass on bone mineral density: the Hordaland Health Study. *Maturitas* 59(2):191–200. <https://doi.org/10.1016/j.maturitas.2007.11.002>
 35. Vinknes KJ, Elshorbagy AK, Nurk E, Dreven CA, Gjesdal CG, Tell GS, Nygard O, Vollset SE, Refsum H (2013) Plasma stearoyl-CoA desaturase indices: association with lifestyle, diet, and body composition. *Obesity* 21(3):E294–E302. <https://doi.org/10.1002/oby.20011>
 36. Benowitz NL, Jacob P III, Ahijevych K, Jarvis MJ, Hall S, LeHouezec J, Hansson A, Lichtenstein E, Henningfield J, Tsoh J, Hurt RD (2002) Biochemical verification of tobacco use and cessation. *Nicotine Tob Res* 4 (2):149–159. <https://doi.org/10.1080/14622200210123581>
 37. Oyen J, Gram Gjesdal C, Nygard OK, Lie SA, Meyer HE, Apalset EM, Ueland PM, Pedersen ER, Middtun O, Vollset SE, Tell GS (2014) Smoking and body fat mass in relation to bone mineral density and hip fracture: the Hordaland Health Study. *PLoS ONE* 9(3):e92882. <https://doi.org/10.1371/journal.pone.0092882>
 38. Choi E, Park Y (2016) The association between the consumption of fish/shellfish and the risk of osteoporosis in men and postmenopausal women aged 50 years or older. *Nutrients* 8(3):113. <https://doi.org/10.3390/nu8030113>
 39. Chan R, Woo J, Leung J (2011) Effects of food groups and dietary nutrients on bone loss in elderly Chinese population. *J Nutr Health Aging* 15(4):287–294
 40. Melaku YA, Gill TK, Adams R, Shi Z (2016) Association between dietary patterns and low bone mineral density among adults aged 50 years and above: findings from the North West Adelaide Health Study (NWAHS). *Br J Nutr* 116(8):1437–1446. <https://doi.org/10.1017/s0007114516003366>
 41. Okubo H, Sasaki S, Horiguchi H, Oguma E, Miyamoto K, Hosoi Y, Kim MK, Kayama F (2006) Dietary patterns associated with bone mineral density in premenopausal Japanese farmwomen. *Am J Clin Nutr* 83(5):1185–1192
 42. Erkkila AT, Sadeghi H, Isanejad M, Mursu J, Tuppurainen M, Kroger H (2017) Associations of Baltic Sea and Mediterranean dietary patterns with bone mineral density in elderly women. *Public Health Nutr*. <https://doi.org/10.1017/s1368980017001793>
 43. Tang BMP, Eslick GD, Nowson C, Smith C, Bensoussan A (2007) Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet* 370(9588):657–666. [https://doi.org/10.1016/S0140-6736\(07\)61342-7](https://doi.org/10.1016/S0140-6736(07)61342-7)
 44. Heaney RP (2007) Effects of protein on the calcium economy. *Int Congr Ser* 1297:191–197. <https://doi.org/10.1016/j.ics.2006.08.025>
 45. Deutz NE, Bauer JM, Barazzoni R, Biolo G, Boirie Y, Bosy-Westphal A, Cederholm T, Cruz-Jentoft A, Krznaric Z, Nair KS, Singer P, Teta D, Tipton K, Calder PC (2014) Protein intake and exercise for optimal muscle function with aging: recommendations from the ESPEN Expert Group. *Clin Nutr* 33(6):929–936. <https://doi.org/10.1016/j.clnu.2014.04.007>
 46. Heaney RP (1998) Excess dietary protein may not adversely affect bone. *J Nutr* 128(6):1054–1057
 47. Allison RJ, Farooq A, Cherif A, Hamilton B, Close GL, Wilson MG (2017) Why don't serum vitamin D concentrations associate with BMD by DXA? A case of being 'bound' to the wrong assay? Implications for vitamin D screening. *Br J Sports Med*. <https://doi.org/10.1136/bjsports-2016-097130>
 48. Lu Z, Chen TC, Zhang A, Persons KS, Kohn N, Berkowitz R, Martinello S, Holick MF (2007) An evaluation of the vitamin D3 content in fish: is the vitamin D content adequate to satisfy the dietary requirement for vitamin D? *J Steroid Biochem Mol Biol* 103(3–5):642–644. <https://doi.org/10.1016/j.jsbmb.2006.12.010>
 49. Lehmann U, Gjessing HR, Hirche F, Mueller-Belecke A, Gudbrandsen OA, Ueland PM, Mellgren G, Lauritzen L, Lindqvist H, Hansen AL, Erkkila AT, Pot GK, Stangl GI, Dierkes J (2015) Efficacy of fish intake on vitamin D status: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 102(4):837–847. <https://doi.org/10.3945/ajcn.114.105395>
 50. Efsa Panel on Dietetic Products, Nutrition and Allergies (2012) Scientific opinion on the tolerable upper intake level of vitamin D. *EFSA J*. <https://doi.org/10.2903/j.efsa.2012.2813>